



RU CLEAR

About Your Medication **Azithromycin**

- ! Swallow the capsules whole on an empty stomach. Do not eat for 1 hour after them.
- ! Do not take indigestion medicines such as Rennie's, Settlers or Gaviscon whilst taking your capsules.
- ! Do not have sex, even with a condom, for the next 7 days whilst the treatment is working.
- ! If you are in sexual relationship with someone, then they will need to be treated as well before you start having sex again, so that they don't pass Chlamydia back to you again.
- ! Sometimes antibiotics can cause stomach upsets and skin rashes. For more information about side effects read the drug information leaflet that comes with your capsules.

If you are worried about how you feel after taking your medication ring the RU Clear office (confidential Helpline) on 0800 046 1303 or ring your GP.

You have been tested for Chlamydia and Gonorrhoea which are common STI's. The following activities may put you at a higher risk of having another less common infection. These are:

- Having sex with men who have sex with other men (or for females having sex with bi-sexual men)
- Injecting drugs (or being the sexual partner of someone who is an injecting drug user.)
- Having sex with somebody who comes from a country, or has had sex in a country, where HIV is common – particularly places in Southern Africa, South East Asia and Russia.

If you think any of these applies to you we would advise further testing at a GUM clinic

EVERY TIME YOU HAVE SEX WITHOUT A CONDOM, WITH A NEW PARTNER, GET A CHLAMYDIA TEST. IT'S BEST TO ALWAYS USE A CONDOM.

Website Address - www.ruclear.co.uk



RU CLEAR

About Your Medication **Azithromycin**

- ! Swallow the capsules whole on an empty stomach. Do not eat for 1 hour after them.
- ! Do not take indigestion medicines such as Rennie's, Settlers or Gaviscon whilst taking your capsules.
- ! Do not have sex, even with a condom, for the next 7 days whilst the treatment is working.
- ! If you are in sexual relationship with someone, then they will need to be treated as well before you start having sex again, so that they don't pass Chlamydia back to you again.
- ! Sometimes antibiotics can cause stomach upsets and skin rashes. For more information about side effects read the drug information leaflet that comes with your capsules.

If you are worried about how you feel after taking your medication ring the RU Clear office (confidential Helpline) on 0800 046 1303 or ring your GP.

You have been tested for Chlamydia and Gonorrhoea which are common STI's. The following activities may put you at a higher risk of having another less common infection. These are:

- Having sex with men who have sex with other men (or for females having sex with bi-sexual men)
- Injecting drugs (or being the sexual partner of someone who is an injecting drug user.)
- Having sex with somebody who comes from a country, or has had sex in a country, where HIV is common – particularly places in Southern Africa, South East Asia and Russia.

If you think any of these applies to you we would advise further testing at a GUM clinic

EVERY TIME YOU HAVE SEX WITHOUT A CONDOM, WITH A NEW PARTNER, GET A CHLAMYDIA TEST. IT'S BEST TO ALWAYS USE A CONDOM.

Website Address - www.ruclear.co.uk