



About Your Medication Erythromycin

- Swallow your capsule whole with plenty of water during a meal.
- Take at regular intervals and always complete the course.
- Do not take indigestion medicines such as Rennies, Settlers or Gaviscon whilst taking your capsules.
- It is very important that you do not have sex, even with a condom, with anyone whilst you are taking your antibiotics.
- If you are in sexual relationship with someone, then they will need to be treated as well before you start having sex again, so that they don't pass Chlamydia back to you again.
- Sometimes antibiotics can cause stomach upsets and skin rashes. For more information about side effects read the drug information leaflet that comes with your capsules.

If you are worried about how you feel after taking your medication ring the RU Clear office (confidential Helpline) on 0800 046 1303 or ring your GP

You have tested for Chlamydia and Gonorrhoea which are common STI's. The following activities may put you at a higher risk of having another less common infection.

These are:

- Having sex with men who have sex with other men (or for females having sex with bi-sexual men)
- Injecting drugs (or being the sexual partner of someone who is an injecting drug user.)
- Having sex with somebody who comes from a country, or has had sex in a country, where HIV is common – particularly places in Southern Africa, South East Asia and Russia.

If you think any of these applies to you we would advise further testing at a GUM clinic.

ALWAYS GET A TEST BEFORE HAVING SEX WITHOUT A CONDOM WITH A NEW PARTNER. IT'S ALWAYS BEST TO USE A CONDOM.



About Your Medication Erythromycin

- Swallow your capsule whole with plenty of water during a meal.
- Take at regular intervals and always complete the course.
- Do not take indigestion medicines such as Rennies, Settlers or Gaviscon whilst taking your capsules.
- It is very important that you do not have sex, even with a condom, with anyone whilst you are taking your antibiotics.
- If you are in sexual relationship with someone, then they will need to be treated as well before you start having sex again, so that they don't pass Chlamydia back to you again.
- Sometimes antibiotics can cause stomach upsets and skin rashes. For more information about side effects read the drug information leaflet that comes with your capsules.

If you are worried about how you feel after taking your medication ring the RU Clear office (confidential Helpline) on 0800 046 1303 or ring your GP

You have tested for Chlamydia and Gonorrhoea which are common STI's. The following activities may put you at a higher risk of having another less common infection.

These are:

- Having sex with men who have sex with other men (or for females having sex with bi-sexual men)
- Injecting drugs (or being the sexual partner of someone who is an injecting drug user.)
- Having sex with somebody who comes from a country, or has had sex in a country, where HIV is common – particularly places in Southern Africa, South East Asia and Russia.

If you think any of these applies to you we would advise further testing at a GUM clinic.

ALWAYS GET A TEST BEFORE HAVING SEX WITHOUT A CONDOM WITH A NEW PARTNER. IT'S ALWAYS BEST TO USE A CONDOM.